

A NOTE ON INTERMEDIATE SUBFACTORS OF KRISHNAN-SUNDER SUBFACTORS

BINA BHATTACHARYYA

ABSTRACT. A Krishnan-Sunder subfactor $R_U \subset R$ of index k^2 is constructed from a permutation biunitary matrix $U \in M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$, i.e. the entries of U are either 0 or 1 and both U and its block transpose are unitary. The author previously showed that every irreducible Krishnan-Sunder subfactor has an intermediate subfactor by exhibiting the associated Bisch projection. The author has also shown in a separate paper that the principal and dual graphs of the intermediate subfactor are the same as those of the subfactor $R^\Gamma \subset R^H$, where $H \subset \Gamma$ is an inclusion of finite groups with an outer action on R . In this paper we give a direct proof that the intermediate subfactor is isomorphic to $R^\Gamma \subset R^H$.

1. BACKGROUND AND INTRODUCTION

There is a well-known way of constructing subfactors of the hyperfinite II_1 subfactor R from certain squares of finite-dimensional C^* -algebras. Since we will use this construction repeatedly, we review it briefly. Suppose we have a square of finite-dimensional C^* -algebras,

$$(1) \quad \begin{array}{ccc} B_0 & \subset & B_1 \\ \cup & & \cup \\ A_0 & \subset & A_1 \end{array}$$

along with a nondegenerate trace on B_1 . Given any inclusion of algebras $A \subset B$ with a nondegenerate trace on B , let E_A^B denote the unique trace preserving conditional expectations from B to A . The square (1) is a *commuting* square if $E_{A_1}^{B_1}(B_0) = A_0$. The square is *symmetric* if B_1 is linearly spanned by $B_0 A_1$. There are many equivalent conditions to these; see [7], [8] for details. An inclusion of finite-dimensional C^* -algebras $A \subset B$ is *connected* if its Bratteli diagram is connected (equivalently, the centers of B and A have trivial intersection). Assume (1) is a symmetric commuting square with connected inclusions, and the trace is the unique Markov trace of the inclusion $B_0 \subset B_1$. We can construct a ladder of symmetric commuting squares:

$$(2) \quad \begin{array}{ccccccccccc} B_0 & \subset & B_1 & \subset & B_2 & \subset & \cdots & \subset & B_n & \subset & \cdots \\ \cup & & \cup & & \cup & & \cdots & & \cup & & \cdots \\ A_0 & \subset & A_1 & \subset & A_2 & \subset & \cdots & \subset & A_n & \subset & \cdots \end{array}$$

by iterating the basic construction to the right, i.e. do the basic construction to the right on the top row and adjoin the Jones projection to the bottom row (see [8], [7], [10]). Then we may complete the inclusion of algebras $\cup_n A_n \subset \cup_n B_n$ with respect to the unique trace on $\cup_n B_n$ to obtain a hyperfinite II_1 subfactor ([8]).

Fix integers k and p . We will consider squares of the form (4). Index the rows and columns of matrices in $M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$ by the set $\{1, 2, \dots, p\} \times \{1, 2, \dots, k\}$ in the natural way. Following the notation in [9], we denote elements of $\{1, 2, \dots, p\}$ with Greek letters and elements of $\{1, 2, \dots, k\}$ with Roman letters. For F in $M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$, denote the entry of F in row (α, a) and column (β, b) by $F_{\alpha a}^{\beta b}$.

Let U be a permutation matrix in $M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$, i.e. its entries are either 0 or 1. U is a permutation biunitary if the block transpose $\tilde{U}$ of U , defined by

$$(3) \quad \tilde{U}_{\beta a}^{\alpha b} = U_{\alpha a}^{\beta b},$$

is also a permutation matrix. Equivalently, U is a permutation biunitary if

$$(4) \quad \begin{array}{ccc} U(1 \otimes M_k(\mathbb{C}))U^* & \subset & M_p(\mathbb{C}) \otimes M_k(\mathbb{C}) \\ \cup & & \cup \\ \mathbb{C} & \subset & M_p(\mathbb{C}) \otimes 1 \end{array}$$

is a commuting square ([8],[6]). We may then construct a subfactor as described above, which we denote $R_U \subset R$. In [9], Krishnan and Sunder list all the nonequivalent subfactors of this type with $k = p = 3$ and compute the principal graphs of all the finite depth ones.

2. THE INTERMEDIATE SUBFACTOR

In Proposition 2.2, we show that if U is a permutation biunitary then the commuting square (4) may be decomposed into two adjacent symmetric commuting squares,

$$(5) \quad \begin{array}{ccc} U(1 \otimes M_k(\mathbb{C}))U^* & \subset & M_p(\mathbb{C}) \otimes M_k(\mathbb{C}) \\ \cup & & \cup \\ U(1 \otimes \Delta_k)U^* & \subset & M_p(\mathbb{C}) \otimes \Delta_k \\ \cup & & \cup \\ \mathbb{C} & \subset & M_p(\mathbb{C}) \otimes 1 \end{array}$$

where $\Delta_k \cong \mathbb{C}^k$ is the diagonal subalgebra of $M_k(\mathbb{C})$.

An essential ingredient in our analysis of Krishnan-Sunder subfactors is the following result of [9].

Lemma 2.1 (Krishnan-Sunder). *If U is a permutation biunitary in $M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$, then there exist permutations $\lambda_1, \lambda_2, \dots, \lambda_p$ in S_k and permutations $\rho_1, \rho_2, \dots, \rho_k$ in S_p , such that*

$$U_{\alpha a}^{\beta b} = \delta_{\beta, \rho_a(\alpha)} \delta_{b, \lambda_\alpha(a)}$$

Consequently, there are permutations $\nu_1, \nu_2, \dots, \nu_p$ in S_k and permutations $\theta_1, \theta_2, \dots, \theta_k$ in S_p , such that

$$(U^*)_{\alpha a}^{\beta b} = \delta_{\beta, \theta_a(\alpha)} \delta_{b, \nu_\alpha(a)}$$

(In Krishnan and Sunder's notation [9], $\nu = \psi^{-1}$ and $\theta = \phi^{-1}$).

Proposition 2.2. *If U is a permutation biunitary in $M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$, then $U(1 \otimes \Delta_k)U^* \subset M_p(\mathbb{C}) \otimes \Delta_k$, and the two small squares in (5) are symmetric commuting squares.*

Proof. To simplify notation, in this proof we denote $\lambda_\alpha(a)$ and $\rho_a(\alpha)$ in Lemma 2.1 by $\alpha(a)$ and $a(\alpha)$, respectively. Let $\{e_{\alpha, \beta}\}$ and $\{f_{a, b}\}$ be the natural sets of matrix units for $M_p(\mathbb{C})$ and $M_k(\mathbb{C})$, respectively. Then $\{1 \otimes f_{a, a}\}_a$ is a basis of $1 \otimes \Delta_k$, and

$$(6) \quad U(1 \otimes f_{a, a})U^* = U \left(\sum_{\alpha=1}^p e_{\alpha, \alpha} \otimes f_{a, a} \right) U^* = \sum_{\alpha=1}^p e_{a(\alpha), a(\alpha)} \otimes f_{\alpha(a), \alpha(a)}$$

which is contained in $M_p(\mathbb{C}) \otimes \Delta_k$. Hence, $U(1 \otimes \Delta_k)U^* \subset M_p(\mathbb{C}) \otimes \Delta_k$.

Since (4) is a symmetric commuting square, it suffices to prove that the upper square is commuting and the lower square is symmetric.

The trace-preserving conditional expectation E from $M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$ to $M_p(\mathbb{C}) \otimes \Delta_k$, is given by

$$E(e_{\alpha,\beta} \otimes f_{a,b}) = \delta_{a,b}(e_{\alpha,\beta} \otimes f_{a,a})$$

So,

$$\begin{aligned} E(U(1 \otimes f_{a,b})U^*) &= E\left(\sum_{\alpha=1}^p e_{a(\alpha),b(\alpha)} \otimes f_{\alpha(a),\alpha(b)}\right) \\ &= \delta_{a,b} \sum_{\alpha=1}^p (e_{a(\alpha),a(\alpha)} \otimes f_{\alpha(a),\alpha(a)}) \\ &= \delta_{a,b} U(1 \otimes f_{a,a})U^* \end{aligned}$$

Therefore, the upper square of (5) is commuting.

It remains to prove that the lower square is symmetric. For any $b \in \{1, 2, \dots, k\}$ and $\beta \in \{1, 2, \dots, p\}$ there exists a, α such that $U_{\alpha,a}^{\beta,b} = 1$, because U is a permutation matrix. By (6) and the fact that ρ_a is a permutation,

$$e_{a(\alpha),\gamma} \otimes f_{\alpha(a),\alpha(a)} \in (U(1 \otimes \Delta_k)U^*)(M_p(\mathbb{C}) \otimes 1)$$

for all $\gamma \in \{1, 2, \dots, p\}$. Since $\beta = a(\alpha)$ and $b = \alpha(a)$ were chosen arbitrarily, $M_p(\mathbb{C}) \otimes \Delta_k$ is linearly spanned by $(U(1 \otimes \Delta_k)U^*)(M_p(\mathbb{C}) \otimes 1)$. $\square$

Note that the inclusion $U(1 \otimes \Delta_k)U^* \subset M_p(\mathbb{C}) \otimes \Delta_k$ is not necessarily connected. Proposition 2.2 immediately implies,

Corollary 2.3. *If U is a permutation biunitary in $M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$, then $R_U \subset R$ has an intermediate von Neumann subalgebra $R_U \subset P_U \subset R$, i.e. the subalgebra constructed from the lower symmetric commuting square in (5). In particular, if $R_U \subset R$ is irreducible then $R_U \subset P_U \subset R$ is an intermediate subfactor.*

Recall $\tilde{U}$ in (3). We will show in Proposition 2.5 that $R_{\tilde{U}} \subset P_{\tilde{U}}$ and $P_U \subset R$ are dual (by symmetry so are $R_U \subset P_U$ and $P_{\tilde{U}} \subset R$) and can be constructed from a biunitary permutation matrix $\Lambda \in \Delta_p \otimes M_k(\mathbb{C})$

Let the permutations $\lambda_\alpha, \rho_a, \nu_\alpha$, and θ_a be as in Lemma 2.1. Define permutation matrices $\Lambda \in \Delta_p \otimes M_k(\mathbb{C})$ and $P \in M_p(\mathbb{C}) \otimes \Delta_k$ by:

$$\Lambda_{\alpha a}^{\beta b} = \delta_{\beta,\alpha} \delta_{b,\lambda_\alpha(a)} \quad P_{\alpha a}^{\beta b} = \delta_{\beta,\rho_a(\alpha)} \delta_{b,a}$$

and permutation matrices $N \in \Delta_p \otimes M_k(\mathbb{C})$ and $\Theta \in M_p(\mathbb{C}) \otimes \Delta_k$ by:

$$N_{\alpha a}^{\beta b} = \delta_{\beta,\alpha} \delta_{b,\nu_\alpha(a)} \quad \Theta_{\alpha a}^{\beta b} = \delta_{\beta,\theta_a(\alpha)} \delta_{b,a}.$$

It is easy to check that

$$(7) \quad U = \Theta^* \Lambda = N^* P$$

and

$$(8) \quad \tilde{U} = \Theta N^* = \Lambda P^*$$

The following lemma shows that we may replace U in the commuting squares that engender $R_U \subset P_U$ and $P_U \subset R$, by the simpler unitaries N^* and Λ .

Lemma 2.4. *Let $\Lambda, N \in \Delta_p \otimes M_k(\mathbb{C})$ and $P, \Theta \in M_p(\mathbb{C}) \otimes \Delta_k$ be permutation matrices satisfying (7) and (8). Then*

$$(9) \quad \begin{array}{ccccccc} U(1 \otimes \Delta_k)U^* & \subset & M_p(\mathbb{C}) \otimes \Delta_k & = & N^*(1 \otimes \Delta_k)N & \subset & M_p(\mathbb{C}) \otimes \Delta_k \\ \cup & & \cup & & \cup & & \cup \\ \mathbb{C} & \subset & M_p(\mathbb{C}) \otimes 1 & & \mathbb{C} & \subset & M_p(\mathbb{C}) \otimes 1 \end{array}$$

and

$$(10) \quad \begin{array}{ccccccc} U(1 \otimes M_k(\mathbb{C}))U^* & \subset & M_p(\mathbb{C}) \otimes M_k(\mathbb{C}) & \cong & \Lambda(1 \otimes M_k(\mathbb{C}))\Lambda^* & \subset & M_p(\mathbb{C}) \otimes M_k(\mathbb{C}) \\ \cup & & \cup & & \cup & & \cup \\ U(1 \otimes \Delta_k)U^* & \subset & M_p(\mathbb{C}) \otimes \Delta_k & & \Lambda(1 \otimes \Delta_k)\Lambda^* & \subset & M_p(\mathbb{C}) \otimes \Delta_k \end{array}$$

Proof. Note that conjugation by any permutation matrix in $M_p(\mathbb{C}) \otimes \Delta_k$ (such as P) stabilizes $1 \otimes \Delta_k$. So substituting N^*P for U yields (9). Similarly, substitute $U = \Theta^*\Lambda$ in left-hand side of (10), and then conjugate the entire square by Θ to obtain the right-hand side. $\square$

Proposition 2.5. *$P_U \subset R$ and $R_{\tilde{U}} \subset P_{\tilde{U}}$ are dual inclusions.*

Proof. Clearly $\tilde{\tilde{U}} = U$. So by symmetry, Lemma 2.4 implies

$$(11) \quad \begin{array}{ccccccc} \tilde{U}(1 \otimes \Delta_k)\tilde{U}^* & \subset & M_p(\mathbb{C}) \otimes \Delta_k & \cong & \Lambda(1 \otimes \Delta_k)\Lambda^* & \subset & M_p(\mathbb{C}) \otimes \Delta_k \\ \cup & & \cup & & \cup & & \cup \\ \mathbb{C} & \subset & M_p(\mathbb{C}) \otimes 1 & & \mathbb{C} & \subset & M_p(\mathbb{C}) \otimes 1 \end{array}$$

Let $e \in M_k(\mathbb{C})$ be the usual Jones projection of the Jones extension $\mathbb{C} \subset \Delta_k \subset M_k(\mathbb{C})$. Conjugation by Λ fixes $1 \otimes e$, hence (10) is the upward basic construction of (11). Since both squares are symmetric, doing the basic construction to the right yields dual inclusions. $\square$

3. THE SUBGROUP SUBFACTOR

Fix a permutation biunitary U . Let ν_α , $1 \leq \alpha \leq p$, and θ_a , $1 \leq a \leq k$, be the permutations defined before Lemma 2.4.

Let Γ be the subgroup of S_k generated by elements of the form $\nu_\alpha \nu_\beta^{-1}$, i.e.,

$$\Gamma = \Gamma_U = \langle \nu_\alpha \nu_\beta^{-1} : \alpha, \beta \in \{1, 2, \dots, p\} \rangle$$

For each $a \in \{1, 2, \dots, k\}$ let $H_a \subset \Gamma$ be the subgroup that fixes a . If Γ acts transitively on $\{1, 2, \dots, k\}$, then the subgroups H_a are all conjugate. In this case set $H = H_1$. In general, let Ω be the set of orbits in $\{1, 2, \dots, k\}$, and for each $r \in \Omega$ set $H_r = H_a$ for an arbitrary representative a in r .

Remark 3.1. *Krishnan and Sunder use the group $\langle \nu_\alpha^{-1} \nu_\beta : \alpha, \beta \in \{1, 2, \dots, p\} \rangle$ instead of Γ in [9]. However, the two groups, as well as their fixed point subgroups, are equivalent via conjugation by ν_γ for any $\gamma \in \{1, 2, \dots, p\}$. We depart from [9] for notational convenience in the proof of the following theorem.*

Theorem 3.2. *Let Γ act on R by outer automorphisms. There is a canonical isomorphism of $Z(P_U)$ with $\mathbb{C}^\Omega$. If q_r is the minimal projection in $Z(P_U)$ corresponding to $r \in \Omega$, then $q_r R_U \subset q_r P_U$ is isomorphic to $R^\Gamma \subset R^{H_r}$. In particular, if Γ acts transitively on $\{1, 2, \dots, k\}$ then $R_U \subset P_U$ is a subfactor, and $R_U \subset P_U$ is isomorphic to $R^\Gamma \subset R^H$.*

Proof. By construction of Γ , the cosets $\nu_\alpha^{-1}\Gamma$ are the same for all $\alpha \in \{1, 2, \dots, p\}$. Denote this coset by Γ' . Given a set S , let Δ_S denote the algebra of functions $S \rightarrow \mathbb{C}$ with pointwise multiplication and the trace $f \mapsto \frac{1}{|S|} \sum_{s \in S} f(s)$. Denote the characteristic function of $s \in S$ by x_s . Define an inclusion map $i: \Delta_\Gamma \rightarrow M_p(\mathbb{C}) \otimes \Delta_{\Gamma'}$ by $i(x_g) = \sum_\alpha e_{\alpha, \alpha} \otimes x_{\nu_\alpha^{-1}g}$. It is straightforward to check that

$$(12) \quad \begin{array}{ccc} \Delta_\Gamma & \subset_i & M_p(\mathbb{C}) \otimes \Delta_{\Gamma'} \\ & \cup & \cup \\ \mathbb{C} & \subset & M_p(\mathbb{C}) \end{array}$$

is a symmetric commuting square with connected inclusions, if we take the trace on $M_p(\mathbb{C}) \otimes \Delta_{\Gamma'}$ to be the product trace.

Then we can construct a hyperfinite II_1 subfactor $B \subset A$ by iterating the basic construction to the right in the usual way.

$$(13) \quad \begin{array}{ccccccccccc} \Delta_\Gamma & \subset^{\mathcal{G}} & M_p(\mathbb{C}) \otimes \Delta_{\Gamma'} & \subset^{\mathcal{G}^t} & A_2 & \subset^{\mathcal{G}} & A_3 & \subset^{\mathcal{G}^t} & \cdots & A \\ \cup & & \cup & & \cup & & \cup & & & \cup \\ \mathbb{C} & \subset & M_p(\mathbb{C}) & \subset & B_2 & \subset & B_3 & \subset & \cdots & B \end{array}$$

Note that the Bratteli diagram $\mathcal{G}$ as marked in (13) is the bipartite graph with even vertices labeled by Γ , odd vertices labeled by Γ' , and an edge for each pair $(g, \alpha) \in \Gamma \times \{1, 2, \dots, p\}$ going from g to $\nu_\alpha^{-1}g$. We denote the reflection of $\mathcal{G}$ by $\mathcal{G}^t$. For each n , $B_n \subset A_n$ is isomorphic to $M_{p^n}(\mathbb{C}) \otimes 1 \subset M_{p^n}(\mathbb{C}) \otimes \Delta_{\tilde{\Gamma}}$, where $\tilde{\Gamma}$ is Γ or Γ' according to whether n is even or odd. We claim that $B \subset A$ is irreducible. By Ocneanu compactness, $B' \cap A = M_p(\mathbb{C})' \cap \Delta_\Gamma = \Delta_\Gamma \cap (1 \otimes \Delta_{\Gamma'})$. Suppose $\sum_{g \in \Gamma} k_g x_g \in 1 \otimes \Delta_{\Gamma'}$, where $k_g \in \mathbb{C}$. Then

$$\sum_{g \in \Gamma} k_g x_g = \sum_{g' \in \Gamma'} \sum_{\nu_\alpha^{-1}g = g'} k_g (f_{\alpha, \alpha} \otimes x_{g'})$$

Since $\sum k_g x_g \in 1 \otimes \Delta_{\Gamma'}$, we must have that k_g is constant over all $g \in \{\nu_\alpha g'\}_\alpha$. Since this holds for all g' , it follows that k_g is constant over all $g \in \Gamma 1 = \Gamma$. Therefore, $\sum k_g x_g \in \mathbb{C}$. This proves the claim.

For each $g \in \Gamma$, let μ_g be the automorphism of $\mathcal{G}$ that maps each vertex $g' \in \Gamma \cup \Gamma'$ to $g'g^{-1}$ and each edge (g', α) to the edge $(g'g^{-1}, \alpha)$. The morphism is well defined since the endpoints of the edge (g', α) are mapped to the endpoints of $(g'g^{-1}, \alpha)$. Moreover, μ_g obviously preserves the trace weights of $\mathcal{G}$. Clearly $g \mapsto \mu_g$ is a group action of G on $\mathcal{G}$. Now extend μ to the chain of Bratteli diagrams of the top row of inclusions in (13). For each n , μ implements an action μ^n of Γ on A_n by trace preserving automorphisms. The family of actions $\{\mu^n: \Gamma \rightarrow \text{Aut}(A_n)\}_n$ are consistent, i.e. $\mu^n|_{A_{n-1}} = \mu^{n-1}$, and thus extend to an action of Γ on A . We denote this action again by μ . Note that the action of Γ on A is outer since $B \subset A$ is irreducible. Therefore $B \subset A$ is isomorphic to $R^\Gamma \subset R$, as in the statement of the theorem.

Let E^Γ be the group averaging maps from A onto the fixed point algebra A^Γ . The action of μ_g on $A_n = M_{p^n}(\mathbb{C}) \otimes \Delta_{\tilde{\Gamma}}$ is given by $\mu_g(F \otimes x'_g) = F \otimes x_{g'g^{-1}}$, hence $E^\Gamma(A_n) = B_n$ for each n . Therefore, $A^\Gamma = B$.

We first assume that Γ acts transitively on $\{1, 2, \dots, k\}$. Define an inclusion $\Delta_{\Gamma/H} \rightarrow \Delta_\Gamma$ by $x_{gH} \mapsto \sum_{g' \in gH} x_{g'}$. Similarly define $\Delta_{\Gamma'/H} \rightarrow \Delta_{\Gamma'}$. For each n , let $C_n = A_n^H$. Clearly $C_n = B_n \otimes \Delta_{\tilde{\Gamma}/H}$

where $\tilde{\Gamma}$ is Γ or Γ' according to whether n is even or odd. Thus we have an intermediate chain of algebras

$$(14) \quad \begin{array}{ccccccccccc} \Delta_{\Gamma} & \subset^{\mathcal{G}} & M_p(\mathbb{C}) \otimes \Delta_{\Gamma'} & \subset^{\mathcal{G}^t} & A_2 & \subset^{\mathcal{G}} & A_3 & \subset^{\mathcal{G}^t} & \cdots & A \\ \cup & & \cup & & \cup & & \cup & & & \cup \\ \Delta_{\Gamma/H} & \subset & M_p(\mathbb{C}) \otimes \Delta_{\Gamma'/H} & \subset & C_2 & \subset & C_3 & \subset & \cdots & C \\ \cup & & \cup & & \cup & & \cup & & & \cup \\ \mathbb{C} & \subset & M_p(\mathbb{C}) & \subset & B_2 & \subset & B_3 & \subset & \cdots & B \end{array}$$

where $C = A^H$. Since the group averaging map E^H from A onto A^H is the conditional expectation from A_n onto C_n for each n , it follows that the upper-left-most square of (14) is commuting. It is straightforward to verify that the lower-left-most square of (14) is symmetric, hence both of the left-most squares are symmetric commuting squares. For $n \geq 2$, C_n contains the Jones projection of the inclusion $A_{n-2} \subset A_{n-1}$, hence the chain $(C_n)_n$ contains the Jones tower of $C_0 \subset C_1$. Moreover, the Bratteli diagram of $C_{n-1} \subset C_n$ is the transpose of $C_{n-2} \subset C_{n-1}$ for $n \geq 2$, hence by dimension considerations, the chain $(C_n)_n$ is no more than the Jones tower. Therefore, both the upper and the lower ladders are the ones obtained by iterating the basic construction in the usual way from the left-most square.

Now consider the lower left square $(*)$ of (14). We claim that $(*)$ is isomorphic to (9) via the identification of Γ/H and Γ'/H with $\{1, 2, \dots, k\}$, by $gH \mapsto g(1)$. Let $(\hat{a})_{a=1}^k$ be the minimal projections in Δ_k , and define an isomorphism $M_p(\mathbb{C}) \otimes \Delta_{\Gamma'/H} \rightarrow M_p(\mathbb{C}) \otimes \Delta_k$ by $F \otimes x_{gH} \mapsto F \otimes \hat{g}(1)$. Fix a and choose $f \in \Gamma$ such that $f(1) = a$. Then $N^*(1 \otimes \hat{a})N = \sum_{\alpha} e_{\alpha, \alpha} \otimes (\nu_{\alpha}^{-1}(a))$ is the image of $\sum_{\alpha} e_{\alpha, \alpha} \otimes x_{\nu_{\alpha}^{-1}fH} = 1 \otimes x_{fH}$. Therefore, $(*)$ is isomorphic to (9), and $R_U \subset P_U$ is isomorphic to $A^{\Gamma} \subset A^H$. This proves $R_U \subset P_U$ is isomorphic to $R^{\Gamma} \subset R^H$, as in the statement of the theorem.

If Γ does not act transitively, then $N^*(1 \otimes \Delta_k)N \subset M_p(\mathbb{C}) \otimes \Delta_k$ is not a connected inclusion; its connected components correspond to the orbits of Γ in $\{1, 2, \dots, k\}$. Given an orbit $r \in \Omega$, let $q_r = N^* \sum_{a \in r} \hat{a}N$. Clearly q_r is central in P_U and $q_r R_U \subset q_r P_U$ can be obtained by iterating the basic construction on

$$\begin{array}{ccc} q_r N^*(1 \otimes \Delta_k)N & \subset & q_r (M_p(\mathbb{C}) \otimes \Delta_k) \\ \cup & & \cup \\ q_r \mathbb{C} & \subset & q_r (M_p(\mathbb{C}) \otimes 1) \end{array}$$

By an identical argument as the one above (using the group H_r instead of H), $q_r R_U \subset q_r P_U$ is isomorphic to $A^{\Gamma} \subset A^{H_r}$. This proves $q_r R_U \subset q_r P_U$ is isomorphic to $R^{\Gamma} \subset R^{H_r}$, as in the statement of the theorem. Then, $q_r R_U \subset q_r P_U$ is a subfactor for each r , which implies that $Z(P_U) = \bigoplus_{r \in \Omega} \mathbb{C} q_r$.

□

Corollary 3.3. *Let $H \subset \Gamma$ be any inclusion of finite groups, and let Γ act on the hyperfinite II_1 factor R by an outer action. Let $k = |\Gamma/H|$. Suppose the action of Γ on Γ/H can be generated by p' elements of Γ . Then there exists a permutation biunitary $U \in \Delta_{p'+1} \otimes M_k(\mathbb{C})$ such that $R_U \subset P_U$ (as defined in Corollary 2.3) is isomorphic to $R^{\Gamma} \subset R^H$.*

Proof. Let $U_1, U_2, \dots, U_{p'}$ be $k \times k$ permutation matrices that generate the action of Γ on Γ/H . Set U_0 to be the identity matrix. Let $\{e_{\alpha}\}_{0 \leq \alpha \leq p'}$ be a basis of $\Delta_{p'+1}$, and set $U = \sum_0^{p'} e_{\alpha} \otimes U_{\alpha}$. Obviously $\Gamma_U = \Gamma$ and the fixed point subgroup of Γ 's action on $\{1, 2, \dots, k\}$ is H . By Theorem 3.2, this U does the job. □

4. THE BISCH PROJECTION

We now show that the Bisch projection defined in [1] (see also [3]) corresponds to the intermediate sub-von Neumann algebra $R_U \subset P_U$.

The upward basic construction of $M_p(\mathbb{C}) \otimes 1 \subset M_p(\mathbb{C}) \otimes M_k(\mathbb{C})$ in (4) is $M_p(\mathbb{C}) \otimes M_{k^2}(\mathbb{C})$. The matrix rows and columns of its subalgebra $1 \otimes M_{k^2}(\mathbb{C})$ are indexed naturally by the set $\{1, 2, \dots, k\} \times \{1, 2, \dots, k\}$ ([6], [8]). Given $x \in 1 \otimes M_{k^2}(\mathbb{C})$, denote by x_{ab}^{cd} the entry of x in row $\{a, b\}$ and column $\{c, d\}$. The first relative commutant of $R_U \subset R$ is the subalgebra of $1 \otimes M_{k^2}(\mathbb{C})$ satisfying the Ocneanu compactness condition [6].

Define $p \in 1 \otimes M_{k^2}(\mathbb{C})$ by

$$(15) \quad p_{ab}^{cd} = \begin{cases} 1, & \text{if } a = b = c = d; \\ 0, & \text{otherwise} \end{cases}$$

Proposition 4.1. *The projection p defined above is contained in the first relative commutant of $R_U \subset R$; and p is a Bisch projection, that is, $pndp$ implements the conditional expectation from R to $\{p\}' \cap R$ with respect to the trace.*

Proof. This is proved in somewhat different notation in Lemma 6.4.1 of [1]. For the convenience of the reader we give a proof here. We first show that p is in the first relative commutant using Jones' diagrammatic formulation of the higher relative commutants of $R'_U \cap R$ [8].

We claim that for $a, b \in \{1, 2, \dots, k\}$ and $\alpha, \beta \in \{1, 2, \dots, p\}$:

$$(16) \quad \begin{array}{ccc} & \begin{array}{c} b \\ \uparrow \\ \beta \end{array} & \begin{array}{c} b' \\ \downarrow \\ \alpha \end{array} \\ & \begin{array}{c} \downarrow \\ a \end{array} & \begin{array}{c} \uparrow \\ a' \end{array} \\ & \beta & \xrightarrow{\quad} \alpha \end{array} = \begin{cases} \delta_{\alpha, \beta} \cdot \delta_{b, b'} \cdot \delta_{a, \nu_\beta(b)}, & \text{if } a = a' \\ \delta_{\alpha, \beta} \cdot \delta_{a, a'} \cdot \delta_{a, \nu_\beta(b)}, & \text{if } b = b' \end{cases}$$

The claim is obvious from Section 5 of [2], but here is a direct proof. The left-hand side of (16) is by definition $\sum_{\gamma \in \{1, 2, \dots, p\}} U_{\gamma a}^{\beta b} \overline{U_{\gamma a'}^{\alpha b'}}$. Note that in our case entries of U are either 0 or 1, so $\overline{U} = U$. We have:

$$\begin{aligned} \sum_{\gamma \in \{1, 2, \dots, p\}} U_{\gamma a}^{\beta b} \overline{U_{\gamma a'}^{\alpha b'}} &= \sum_{\gamma} \delta_{\beta, \rho_a(\gamma)} \delta_{b, \lambda_\gamma(a)} \delta_{\alpha, \rho_{a'}(\gamma)} \delta_{b', \lambda_\gamma(a')} \\ &= \delta_{\alpha, \beta} \delta_{b, b'} \delta_{b, \lambda_{\rho_a^{-1}(\alpha)}(a)} && \text{if } a = a' \\ &= \delta_{\alpha, \beta} \delta_{b, b'} \delta_{a, \nu_\alpha(b)} && \text{(Lemma 5 of [9])} \end{aligned}$$

and

$$\begin{aligned} \sum_{\gamma \in \{1, 2, \dots, p\}} U_{\gamma a}^{\beta b} \overline{U_{\gamma a'}^{\alpha b'}} &= \sum_{\gamma} (U^*)_{\beta b}^{\gamma a} (U^*)_{\alpha b'}^{\gamma a'} \\ &= \sum_{\gamma} \delta_{\gamma, \theta_b(\beta)} \delta_{a, \nu_\beta(b)} \delta_{\gamma, \theta_{b'}(\alpha)} \delta_{a', \nu_\alpha(b')} \\ &= \delta_{\alpha, \beta} \delta_{a, a'} \delta_{a, \nu_\beta(b)} && \text{if } b = b' \end{aligned}$$

This proves the claim. Using (16), it is easy to verify that p satisfies the diagrammatic condition for p to be in $R'_U \cap R_1$ (see Theorem 6.1.4 and the preceding discussion in [8]).

Let $q \in M_{k^2}(\mathbb{C})$ be the projection identified with p , that is, $p = 1 \otimes q \in M_p(\mathbb{C}) \otimes M_{k^2}(\mathbb{C})$. Define $p_n = 1 \otimes 1_{M_{k^n}(\mathbb{C})} \otimes q \in M_p(\mathbb{C}) \otimes M_{k^{n+2}}(\mathbb{C})$, for $n = 0, 1, 2, \dots$. The same argument as above shows that p_n is contained in $(n-1)$ st relative commutant of $R_U \subset R$. It is easy to verify that the sequence of projections (p_n) along with the sequence of Jones projections (e_n) of $R_U \subset R$ satisfies

the Bisch-Jones relations of the Fuss-Catalan algebras [5] with $\alpha = \beta = k$. It follows by [4] that p implements the conditional expectation from R to $\{p\}' \cap R$ with respect to the trace $\square$

Proposition 4.2. *Let p be the Bisch projection defined above and let E_p be the conditional expectation from R to $\{p\}' \cap R$ implemented by p (by Proposition 4.1). Then $P_U = E_p(R)$.*

Proof. Let e be the Jones projection of the extension $R_U \subset R \subset R_1$. Let (A_m) , (C_m) , and (B_m) , $m = 0, 1, 2, \dots$, be the chains of algebras obtained by iterating the basic construction on (5) to the right, and let $D_m = \langle B_m, e \rangle$:

$$\begin{array}{ccccccccccc}
 (17) & D_0 & \subset & D_1 & \subset & D_2 & \subset & D_3 & \subset & \cdots & R_1 \\
 & \cup & & \cup & & \cup & & \cup & & & \cup \\
 & U(1 \otimes M_k(\mathbb{C}))U^* & \subset & M_p(\mathbb{C}) \otimes M_k(\mathbb{C}) & \subset & A_2 & \subset & A_3 & \subset & \cdots & R \\
 & \cup & & \cup & & \cup & & \cup & & & \cup \\
 & U(1 \otimes \Delta_k)U^* & \subset & M_p(\mathbb{C}) \otimes \Delta_k & \subset & C_2 & \subset & C_3 & \subset & \cdots & P_U \\
 & \cup & & \cup & & \cup & & \cup & & & \cup \\
 & \mathbb{C} & \subset & M_p(\mathbb{C}) & \subset & B_2 & \subset & B_3 & \subset & \cdots & R_U
 \end{array}$$

The type of construction above is well-known ([6], also see Ocneanu compactness in [8]), and in particular has the property that the chain $B_m \subset C_m \subset A_m \subset D_m$ has the same Bratteli diagram for all odd m . In other words, $B_m \subset C_m \subset A_m \subset D_m$ is isomorphic to $B_m \subset B_m \otimes \Delta_k \subset B_m \otimes M_k(\mathbb{C}) \subset B_m \otimes M_{k^2}(\mathbb{C})$ for all odd m .

By definition, p is a projection in $1 \otimes M_{k^2}(\mathbb{C}) \subset D_1 = M_p(\mathbb{C}) \otimes M_{k^2}(\mathbb{C})$. Let $q \in M_{k^2}(\mathbb{C})$ be the projection identified with p , that is, $p = 1 \otimes q \in M_p(\mathbb{C}) \otimes M_{k^2}(\mathbb{C})$. Using (15), it is easy to check:

$$(18) \quad q \in (\Delta_k \otimes 1)' \cap M_{k^2}(\mathbb{C})$$

$$(19) \quad q(M_k(\mathbb{C}) \otimes 1)q = (\Delta_k \otimes 1)q.$$

It follows that $p \in C'_1 \cap D_1$ and $pA_1p = C_1p$, hence $C_1 = E_p(A_1)$.

By Proposition 4.1, p is contained in $R'_U \cap R_1$, hence p is *flat* [6]. By flatness, p , as an element of $B_m \otimes M_{k^2}(\mathbb{C}) \cong D_m$, for m odd, is identified with $1_{B_m} \otimes q \in B_m \otimes M_{k^2}(\mathbb{C})$. Recalling that $B_m \subset C_m \subset A_m \subset D_m$ is isomorphic to $B_m \subset B_m \otimes \Delta_k \subset B_m \otimes M_k(\mathbb{C}) \subset B_m \otimes M_{k^2}(\mathbb{C})$, we have by (18) and (19) that $p \in C'_m \cap D_m$ and $pA_mp = C_mp$ for all odd m . Hence, $C_m = E_p(A_m)$ for all odd m . Then by weak continuity of E_p , $P_U = E_p(R)$. $\square$

We restate Proposition 4.2 and Proposition 3.2 as follows:

Corollary 4.3. *The intermediate sub-von Neumann algebra of $R_U \subset R$ corresponding to the Bisch projection p is $R_U \subset P_U$, as defined in Corollary 2.3.*

REFERENCES

- [1] Bina Bhattacharyya, *Krishnan-Sunder subfactors and a new countable family of subfactors related to trees*, Ph.D. thesis, University of California at Berkeley, 1998.
- [2] ———, *Group actions on graphs related to Krishnan-Sunder subfactors*, Trans. Amer. Math. Soc. **355** (2003), no. 2, 433–463.
- [3] Bina Bhattacharyya and Zeph Landau, *Intermediate standard invariants and intermediate planar algebras*, to appear.
- [4] D. Bisch, *A note on intermediate subfactors*, Pacific J. Math. **163** (1994), no. 2, 201–216.
- [5] D. Bisch and V. F. R. Jones, *Algebras associated to intermediate subfactors*, Invent. Math. **128** (1997), no. 1, 89–157.

- [6] A. Ocneanu (Lecture Notes by Y. Kawahigashi), *Quantum symmetry, differential geometry of finite graphs and classification of subfactors*, 1990, University of Tokyo Seminar Notes.
- [7] F. Goodman, P. de la Harpe, and V. F. R. Jones, *Coxeter graphs and towers of algebras*, MSRI Publications, vol. 14, Springer, 1989.
- [8] V. F. R. Jones and V. S. Sunder, *Introduction to subfactors*, London Mathematical Society Lecture Note Series, vol. 234, Cambridge University Press, 1997.
- [9] U. Krishnan and V. S. Sunder, *On biunitary permutation matrices and some subfactors of index 9*, Trans. Amer. Math. Soc. **348** (1996), no. 12, 4691–4736.
- [10] S. Popa, *Orthogonal pairs of $*$ -subalgebras in finite von Neumann algebras*, J. Operator Theory **9** (1983), no. 2, 253–268.

DEEPHAVEN, 4699 OLD IRONSIDES DR., #210, SANTA CLARA, CA 94054, US

E-mail address: `Bina_Bhattacharyya_91@post.harvard.edu`